arXiv:1701.03256v2 [gr-qc] 20 Jan 2017

Topological Quantum Numbers of Dyonic Fields over Taub-Bolt

Daniel Flores-Alfonso, Hernando Quevedo

Instituto de Ciencias Nucleares, Universidad Nacional Autónoma de México, AP 70543, Ciudad de México 04510, México

Abstract

We study the global structure of dyonic fields on a Taub-Bolt background, and compare our results with similar fields over Taub-NUT and Schwarzschild spaces. It is shown that the electric and magnetic charges are quantized and lead to a classification of dyonic Taub-Bolt configurations. The quantization condition for the magnetic charge is similar to that of Dirac's monopole.

Keywords: Topological Charge, Instantons, Taub-Bolt, Dyons

1. Introduction

The study of gravitational instantons in Euclidean Einstein gravity was initiated in the 70's [1, 2, 3, 4, 5, 6] to apply the path-integral approach in the hope to construct a theory of quantum gravity. This expectation was based upon the success reached by this approach in the application of quantum field theory to Yang-Mills theories. Indeed, the instanton solutions of classical Euclidean Yang-Mills field equations, with self-dual field strength, allow one to interpret the physical vacuum state as a superposition of an infinite number of vacuum states. This leads to an alternative non-perturbative quantization of Yang-Mills theories by using the path-integral approach. The idea of applying similar analysis to Euclidean Einstein gravity found several conceptual and technical difficulties [7], which are currently under investigation to formulate a consistent theory of quantum gravity. Nevertheless, gravitational instantons became a topic of intensive research, and together

Email addresses: daniel.flores@correo.nucleares.unam.mx (Daniel Flores-Alfonso), quevedo@nucleares.unam.mx (Hernando Quevedo)

with monopoles and solitons constitute the area of theoretical physics which today is known as topological defects.

A different approach was recently proposed in which concepts of topological quantization are applied to find quantum information from classical fields, including gravitational configurations [8]. In this case, the underlying classical structure is different. To any classical configuration, which includes gauge fields, gravitational fields and mechanical systems, one associates a principal fiber bundle where the base space should contain all the information of the classical configuration and the standard fiber corresponds to its physical symmetries. In the case of gravity, for instance, the base space is the spacetime and the standard fiber is the Lorentz group $SO(3,1)$ so that the bundle is 10-dimensional. The main goal of topological quantization is to show that the topological and geometric properties of the principal fiber bundle can be used to obtain quantum information about the corresponding classical configuration. This goal has been reached only partially [9, 10, 11, 12, 13]. In particular, the topological numbers of the bundle are used to construct a topological spectrum whose physical information is directly related to the information contained in the spectrum obtained in canonical quantization. Topological invariants are therefore an important tool of topological quantization and this is one of the reasons why we are interested in analyzing the properties of topological defects.

Recently, the classic subject of topological defects [14, 15, 16, 17, 18] has been applied to condensed matter, i.e., topological insulators — for a review on the subject see, for example, Ref. [19]. The main rôle is played by topological quantum numbers such as Chern invariants. The subject includes other aspects of topology as well. These subjects are related physically to topologically protected states, time-reversal symmetry, topological phases of matter and topological phase transitions (for a a selection of recent references see [20, 21, 22, 23, 24, 25, 26, 27, 28] and earlier works in [29, 30, 31, 32, 33]). The importance of this research was notoriously recognized in the 2016 Nobel prize which was awarded to D.J. Thouless, F.D.M. Haldane and J.M. Kosterlitz "for theoretical discoveries of topological phase transitions and topological phases of matter".

Phase transitions which include a change in topology have been studied in gravitation as well. Research in this direction usually entails using gravitational instantons — the gravitational analogs of Yang-Mills pseudoparticles[1]. These solutions have positive definite metric. Euclidean metrics may be hard to be interpreted physically, but are a useful tool for calculating

the free energy of the corresponding Lorentzian system. For example, in the presence of a negative cosmological constant, radiation over the AdS space and the AdS-Schwarzschild black hole are phases of the same thermodynamic system. The partition function may be estimated using their Wick rotated counterparts. Topology changes during these phase transitions; thermodynamics [34] and Ricci flow with surgery [35] are used to grasp these changes. These transitions have been studied for Taub-NUT and Taub-Bolt spacetimes as well [36, 37].

It is very well known [5, 38] that a self-dual Maxwell field naturally lives on the Taub-NUT instanton. This field is dyonic. It is a renowned[1] result that the Taub-NUT instanton is topologically trivial. One usually expects that the Chern number of such spaces vanish. However, this is not the case for Taub-Bolt. In this work, we study dyonic Maxwell fields over Taub-Bolt which are not self-dual. We focus on electromagnetic charge, as a topological quantum number, on a Taub-Bolt space — which is a $4-$dimensional gravitational instanton.

Topological quantum numbers originate with the Dirac monopole which has magnetic charge p and gauge potential

$$A = p\cos\theta d\phi \ . \tag{1}$$

Since we are mainly concerned with gravitation in this paper, we review its unique self-gravitating equivalent, i.e., the magnetic Reissner-Nordström spacetime. The gauge potential is still given by Eq.(1), but is defined over the space $\mathbb{R}^2 \times S^2$, the topology of the Reissner-Nordström black hole. In spherical coordinates, A is singular at $\theta = 0$ and $\theta = 2\pi$. To resolve this issue, we consider two gauge potentials equivalent to Eq. (1) each defined on complementary open covers of the spheres coordinated by (θ, ϕ).

$$A_\pm = p(\pm 1 + \cos\theta)d\phi \ . \tag{2}$$

Notice that $A_+ - A_- = 2pd\phi$. Consistency of the $U(1)-$bundle where A is determined requires

$$2p = \text{integer} \ , \tag{3}$$

meaning that the magnetic charge is quantized. Moreover, the topology of the bundle space is then $S^3/\mathbb{Z}_n$ whenever $p = n/2$; hence p is called a topological charge.

Alternatively, we can look at the Chern invariants of the configuration. Since our base manifold is four dimensional, there are two relevant Chern

classes to consider, i.e., the two-form [6]

$$c_1 = -\frac{1}{2\pi}\,\mathrm{Tr}\,F\ , \tag{4}$$

and the four-form

$$c_2 = \frac{1}{8\pi^2}(\mathrm{Tr}\,F\wedge F - \mathrm{Tr}\,F\wedge \mathrm{Tr}\,F)\ . \tag{5}$$

This last cohomology class vanishes for $U(1)-$bundles, the gauge group of electromagnetism, so we consider only the two-form c_1. This Chern class corresponds to the field strength, defined by $F = dA$. The sphere representing the event horizon and all 2D submanifolds that can be deformed into it form a special homology class. This leads to an integral formula

$$\int_{S^2} c_1 = 2p\ , \tag{6}$$

Since the Chern class is an integer, we obtain, once again, Eq.(3).

Another question is what happens if the geometry and field are that of electric Reissner-Nordström; this question has been tackled in [8]. Moreover, the generalization to include rotation has been considered as well. Indeed, the Schwarzschild spacetime may be generalized to Kerr-Newman, but independently may be generalized to spaces with a NUT parameter. Specifically, we consider charged versions of Taub-NUT spaces.

This work is organized as follows. In Sec. 2, we review the main properties of the gravitational instantons that are generated by applying a Wick rotation to the Taub-NUT spacetime with electric and magnetic charge. In Sec. 3, we calculate explicitly the topological quantum numbers of dyonic fields on the Euclidean Taub-NUT and Taub-Bolt backgrounds. Finally, in Sec. 4, we conclude with some remarks about possible future applications of our results.

2. Nuts, bolts and dyons

Some gravitational instantons may be obtained through Wick rotating a Lorentzian metric; this is the case of the Taub-NUT and Taub-Bolt[1] instantons. The Lorentzian Taub-NUT spacetime [39, 40] is defined over the space

[1]Taub-Bolt is the counter part of Taub-NUT which has a surface analog to an event horizon; see section 2.

$S^3 \times \mathbb{R}$, and its metric is given by

$$ds^2 = -V(r)(dt + 2L\cos\theta d\phi)^2 + \frac{dr^2}{V(r)} + (r^2 + L^2)(d\theta^2 + \sin^2\theta d\phi^2) \ , \quad (7)$$

with

$$V(r) = \frac{r^2 - 2Mr - L^2}{r^2 + L^2} \ . \quad (8)$$

The underlying topology with periodic time $t \sim t + 8\pi L$ is chosen to avoid the physical presence of Misner strings [41]. The physical parameters are the (ordinary) gravitoelectric mass M and the NUT charge L also called the gravitomagnetic mass. The Taub-NUT metric represents a gravitational dyon, and is asymptotically locally flat (ALF). Indeed, its curvature in the $SO(3,\mathbb{C})-$representation [42], with a local orthonormal metric, can be expressed as [43]

$$\mathcal{R} = \frac{M + iL}{(r + iL)^3} \begin{pmatrix} -1 & & \\ & -1 & \\ & & 2 \end{pmatrix} \ , \quad (9)$$

which is clearly locally flat, although the global topology at infinity can differ from that of the ordinary Euclidean space.

Taub-NUT metrics belong to the Bianchi IX class, i.e., spacetimes with an $SU(2)$ homogeneity group. Bianchi IX metrics have a portrayal as a sequence of squashed three-spheres. Taub-NUT is a very symmetric case as its isometry group is $SU(2)\times U(1)$, where the $U(1)$ subgroup on the right is a cyclic symmetry between two of the invariant directions of the spatial three-spheres. For this reason Taub-NUT is called a biaxial Bianchi IX spacetime. So the sequence portraying the NUT space has squashed circles Hopf fibered over round two-spheres. This singles out one direction, the one in which circles are fibered over spheres.

A generalization of the Taub-NUT metric that includes Maxwell fields was found in [44]. The line element is still written as in Eq.(7), but the function $V(r)$ instead takes the form

$$V(r) = \frac{r^2 - 2Mr - L^2 + Q^2 + p^2}{r^2 + L^2} \ . \quad (10)$$

The construction of this metric takes advantage of the biaxial symmetry described above. The gauge potential is

$$A = \left(\frac{Qr}{r^2 + L^2} + \frac{p}{2L}\frac{r^2 - L^2}{r^2 + L^2} \right) (dt + 2L\cos\theta d\phi) \ . \quad (11)$$

In the $L \to 0$ limit, we recover the dyonic Reissner-Nordström spacetime. Through a simple gauge transformation $A \to A - p/2L\ dt$ we have the more convenient expression

$$A = p\cos\theta d\phi + \left(\frac{Qr - pL}{r^2 + L^2}\right)(dt + 2L\cos\theta d\phi)\ . \tag{12}$$

As before, the gauge potential is singular whenever $\sin\theta = 0$ as apparently the magnetic part of the field vanishes. We resolve this in the classical manner defining

$$A_\pm = p(\pm 1 + \cos\theta)d\phi + \left(\frac{Qr - pL}{r^2 + L^2}\right)(dt + 2L\cos\theta d\phi)\ , \tag{13}$$

concluding that the value $2p$ is an integer. However, the spacetime topology prevents this magnetic charge to be a topological quantum number.

To obtain the gravitational instantons related to this spacetime, we make a Wick rotation $t \mapsto i\tau$ on the neutral Taub-NUT spacetime. Later on, we consider the charged version. However, as is readily seen, the metric is only a real Riemannian metric if we also apply the transformation $L \mapsto iN$. So two geometrically distinct gravitational instantons arise. Additionally, conical singularities emerge unless we impose certain restrictions which are detailed in the sequel. The two Euclidean metrics have the form

$$ds^2 = V(r)(d\tau + 2N\cos\theta d\phi)^2 + \frac{dr^2}{V(r)} + (r^2 - N^2)(d\theta^2 + \sin^2\theta d\phi^2)\ . \tag{14}$$

The S^3 submanifold degenerates at $r = r_+$ where V is nil — the nature of this degeneration distinguishes the instanton pair. Historically, the first case found was the self-dual Taub-NUT instanton [1], so called because its curvature form is self-dual. We henceforth refer to this instanton systematically as Taub-NUT. In what follows Lorentzian spaces are no longer considered so no ambiguity arises. The second case is called Taub-Bolt [45] and was first written down in Ref. [46]. The Taub-Bolt and Taub-NUT manifolds have different allowed ranges for M, N and r; the appropriate conditions have been worked out in [47].

For Taub-NUT, the three-sphere degenerates into a point, called a nut, and V reduces to

$$V_N = \frac{r - N}{r + N}\ . \tag{15}$$

In this coordinate system there is a coordinate singularity at $r = N$ (called a nut). The global structure is that of the space $\mathbb{R}^4$. So the Euler number and Hirzebruch signature of Taub-NUT are $\chi = 1$ and $\sigma = 0$, respectively.

In the Taub-Bolt case, the three-sphere degenerates into a two-sphere, via the Hopf fibration, which is called a bolt. In this case, $V = V_B$ which is

$$V_B = \frac{(r - N/2)(r - 2N)}{r^2 - N^2} \ . \tag{16}$$

As in the NUT case, the bolt (which is at $r = 2N$) is a regular set, but in this coordinate system there is a coordinate singularity. Here $r = N/2$ is not contained within the range of r; moreover, the minimum value of r, r_+, is always greater than N, a property which is also valid in the generalizations of Taub-Bolt we treat later on. The natural topology is that of the space $\mathbb{CP}^2\backslash\{*\}$, which has $\chi = 2$ and $\sigma = 1$.

Both Taub-NUT and Taub-Bolt are limits of compact manifolds with an S^3 boundary. The limit involves taking the boundary to infinity. The bolt in the latter case is a minimal surface very similar to a black hole horizon. Moreover, the space $\mathbb{CP}^2\backslash\{*\}$ has the homotopy type of S^2 which happens in some black holes as well, e.g., Schwarzschild. The Taub-Bolt space has the topology of a cone Hopf fibered over S^2 with the tip of the cone lying on the bolt. Since the cone is retractable, it is straightforward to see that the Taub-Bolt space indeed has the homotopy type of a sphere. Physically, one can think of Taub-Bolt as an excitation of Taub-NUT, just as we can do for Schwarzschild and flat space [45].

As expected, these vacuum solutions may be generalized to gravitational instantons of Euclidean Einstein-Maxwell theory by Wick rotating the dyon potential in Eq. (11). To obtain a real gauge potential, we must take $Q \mapsto iq$. Notice that the quantities L and Q are related to timelike directions in the Lorentzian configuration so Wick rotations affect them. This happens for the angular momentum in rotating configurations as well. Since the parameter p is magnetic, it remains unaffected.

The line element remains with the same structure, but the metric function V instead takes the form

$$V_d = \frac{r^2 - 2Mr + N^2 + p^2 - q^2}{r^2 - N^2} \ , \tag{17}$$

and the dyon potential is

$$A = h(r)(d\tau + 2N\cos\theta d\phi) \ , \tag{18}$$

where

$$h(r) = \frac{qr}{r^2 - N^2} + \frac{p}{2N}\frac{r^2 + N^2}{r^2 - N^2} \ . \tag{19}$$

For these dyonic configurations, consistency again restricts the ranges of the physical parameters, including the electric charge. Given that we require $V_d(r_+) = 0$ and $h(r_+) = 0$, then mass and electric charge must comply with

$$M = \frac{r_+^2 + N^2 + p^2 - q^2}{2r_+} \tag{20}$$

and

$$q = -p\frac{r_+^2 + N^2}{2Nr_+} \ . \tag{21}$$

Since the NUT and Bolt geometry have the same asymptote, the charge and potential at infinity are equal for both cases. The asymptotic charge and potential are q and $p/2N$, respectively.

Taub-NUT and Taub-Bolt configurations have been also studied in generalized theories of gravity [48, 49].

3. Topological quantum numbers

Topological quantum numbers are at the heart of the study of topological defects. They represent the quantized conductance in the quantum Hall effect[29]. They describe the number of levels in molecular spectra[50, 51]. They are the numbers characterizing monopole and instanton solutions[14, 15, 16, 17, 18]. In Skyrmions they represent the baryon number[52]. In $\mathbb{Z}_2$ topological insulators they distinguish between an ordinary insulator state and a quantum Hall state[20].

As a first example, let us consider again the magnetic Reissner-Nordström configuration. The background is a 4D manifold M and has a special 2D submanifold, the two-sphere representing the event horizon[2]. The quantity

$$C_1 = c_1[S^2] = 2p = n \ , \tag{22}$$

is an integer but, it is not a Chern number for M. Chern numbers are obtained from maximally dimensional forms in the Chern classes.

[2]There are infinitely many equivalent special submanifolds, for example the two-sphere at infinity

For an arbitrary unitary group bundle over a 4−dimensional manifold M, the Chern numbers are defined by [6]

$$C_1^2 = c_1 \wedge c_1[M] \ , \qquad C_2 = c_2[M] \ , \tag{23}$$

where

$$c_1 = -\frac{1}{2\pi} \operatorname{Tr} F \ , \qquad c_2 = \frac{1}{8\pi^2}(\operatorname{Tr} F \wedge F - \operatorname{Tr} F \wedge \operatorname{Tr} F) \ . \tag{24}$$

Since $SU(2)$ has traceless field strength, then the number C_1^2 is zero and C_2 simplifies to the usual expression. While for $U(1)$ the inverse happens, i.e., $C_2 = 0$ and

$$C_1^2 = \frac{1}{4\pi^2} \int_M F \wedge F \ . \tag{25}$$

So we readily see that for magnetic Reissner-Nordström $C_1^2 = 0$. For this configuration all Chern numbers vanish, this points to a triviality. It does so in the following sense. The configuration is basically the Dirac monopole multiplied trivially with the space $\mathbb{R}^2$. The fact that the construction is trivial is reflected in the vanishing of the topological numbers.

The manifolds underlying Taub-NUT/Bolt both have a 3−dimensional boundary, S^3_∞ say. Consequently, the Chern-number may also be calculated through the Chern-Simons 3-form integrated on this boundary, i.e.

$$c_1 \wedge c_1[M] = cs[\partial M] = \frac{1}{4\pi^2} \int_{S^3_\infty} A \wedge dA \ , \tag{26}$$

where we have simplified the Chern-Simons form for the Abelian case.

Before continuing to the next section let us consider a Reissner-Nordström instanton with electric charge q and magnetic charge p. Imposing self-duality on the field yields [53]

$$F = \frac{p}{r^2} \left(dr \wedge d\tau - r^2 \sin\theta d\theta \wedge d\phi \right) \ . \tag{27}$$

This field has been studied in [54] and from there we know that the configuration is characterized by

$$C_1 = n \quad \text{and} \quad C_1^2 = 2n^2 \ , \tag{28}$$

where n is an integer and $2p = n$. Note that self-duality also affects the background geometry. Since the metric parameters comply with $p^2 = q^2$ then the background is in fact Schwarzschild. This is a trivial solution to the Euclidean Einstein-Maxwell equations. Since the energy-momentum tensor vanishes and so the background is a solution to the vacuum equations.

Recall that the Schwarzschild instanton is asymptotically flat. Moreover, if we consider the same field as described above over a globally flat space, it is still a solution to the Euclidean Einstein-Maxwell equations albeit a trivial one. This pair of configurations has the same asymptotic boundary and the same field content. The integrals involved in the calculation of Chern numbers yield identical results. The key difference is the cohomology of each background. While it is trivial for flat space it is not for the Schwarzschild instanton. In fact, the second cohomology group of the Schwarzschild space is infinite cyclic i.e. $\mathbb{Z}$. This means that $2p = n$ only for the Schwarzschild background.

3.1. Dyonic Taub-Bolt

In this section, we compare dyonic and self-dual fields over Taub-NUT and Taub-Bolt spaces. To this end, we first review some known results for the Taub-NUT space [53] some of which carry over to the Taub-Bolt space. Firstly, recall that the biaxial symmetry present in the Taub spaces is $U(1)$. The Killing form associated to this symmetry, ξ say, may be used to define a $U(1)$ two-form $F = \lambda d\xi$, where λ is related to the electromagnetic charges. In turn, F has the (anti) self-dual components

$$F_{\pm} = \frac{\lambda(M \pm N)}{(r \pm N)^2} \left[dr \wedge (d\tau + 2N \cos\theta d\phi) \mp (r^2 - N^2) \sin\theta d\theta \wedge d\phi \right] \ . \quad (29)$$

This should be compared with the dyon field whose (anti) self-dual components are obtained from the above expression by changing

$$\lambda(M \pm N) \to 1/2(\pm p - q) \ . \quad (30)$$

When $M = N$, the famous [53, 5, 38, 54] self-dual field strength (on Taub-NUT) is recovered

$$F = \frac{p}{(r+N)^2} \left[dr \wedge (d\tau + 2N \cos\theta d\phi) - (r^2 - N^2) \sin\theta d\theta \wedge d\phi \right] \ . \quad (31)$$

Since the particular case $M = -N$ is not physical, then no anti self-dual field is possible over the Taub-NUT space. Additionally, note that both (anti)

self-dual fields comply with $p^2 - q^2 = 0$. This means that the dyon over a Taub-NUT space does not modify the background. This is a particular distinction between (anti) self-dual fields over Riemannian and Lorentzian backgrounds, as pointed out in [55]. The energy-momentum tensor vanishes for fields with this symmetry and so vacuum solutions may be endowed with them without further changes. This is to say, they are trivial solutions to the Euclidean Einstein-Maxwell equations.

For the dyon over the Taub-Bolt space this is not so. The dyon field is not (anti) self-dual and so the charged Taub-Bolt geometry is different from its neutral counterpart. As mentioned above, consistency requires $V_d(r_+) = 0$ and $h(r_+) = 0$. In this case, $r_+ = r_b$, the only physical solution ($r_b > N$) which satisfies the cubic equation

$$r_+^3 - 2N(1+\nu^2)r_+^2 + 2\nu^2 N^3 = 0 \ , \tag{32}$$

where ν is short hand for $p/2N$.

We recall that the asymptotic behavior of Taub-NUT and Taub-Bolt spaces is identical. For example, the total magnetic flux[3] is

$$\Phi = \frac{1}{2\pi} \int\limits_{S^2_\infty} F = -2p \ . \tag{33}$$

However, the cohomology of Taub-NUT is trivial and so we may not conclude that this flux is quantized in general. This means that all continuous values of p are admissible and since F is self-dual, it corresponds to a continuous family of instantons with finite action [54]. The situation is completely different in the case of the Taub-Bolt space since its second cohomology group is infinite cyclic, $H^2(\mathbb{CP}^2 \backslash \{*\}) = \mathbb{Z}$.

The Chern number is calculated as

$$\frac{1}{4\pi^2} \int F \wedge F = 4p^2 \ , \tag{34}$$

where the domain of the integral is not indicated because the result is the same for Taub-NUT ($\mathbb{R}^4$) and Taub-Bolt ($\mathbb{CP}^2 \backslash \{*\}$). This is due to the fact that both dyons vanish at the source (i.e. nut or bolt, depending on the case) and both have the same value at infinity. An alternative way to see this is

[3]For comparison with Chern invariants we normalize the flux with a $1/2\pi$ factor.

to use the fact that both spaces have the same asymptotic boundary and, therefore, the Chern-Simons invariant in both cases is

$$\frac{1}{4\pi^2}\int\limits_{S^3_\infty} A\wedge dA = 4p^2\ . \tag{35}$$

Nevertheless, this number is an integer only for the Taub-Bolt space. In fact, this invariant classifies the different possible gauge potentials — different according to their magnetic charge. This may be paraphrased to saying that the magnetic charge is a topological quantum number for Taub-Bolt. For Taub-NUT, however, the continuous family of fields are referred to as non-topological [54], since the above quantities are non-zero but are not predictable from the background topology.

As a final remark we mention that for $U(1)$ bundles, S^2 backgrounds are of interest — as we know from the Dirac monopole. Since bundle theory is homotopy invariant[4], the same applies for manifolds of higher dimension but with the homotopy type of a two-sphere, e.g., the Taub-Bolt space. Integral cohomology can be forced over the NUT space [53] however, it emerges naturally on the Bolt scenario. Chern classes have already been studied for (anti)self-dual fields over rotating Taub-Bolt spaces [55]. However, to the best of our knowledge the topological quantum numbers of Taub-Bolt dyons (which are not self-dual) have not been studied in the literature before.

4. Conclusions

In this work, we have analyzed the gravitational instantons that arise from the Taub-NUT spacetime with electric and magnetic charges. In particular, we investigated the topological properties of dyons and (anti) self-dual fields on Euclidean backgrounds corresponding to Taub-NUT and Taub-Bolt spaces. We conclude that instantons with a bolt carry topological quantum numbers. Their cohomology is integral and there are countably infinite distinct $U(1)$-bundles over them. We have shown that the Chern numbers of Taub-Bolt dyons are closely related to the invariants of the famous Taub-NUT self-dual field. In terms of physical parameters they are the same. The essential difference is that the former are quantized due to topological reasons.

[4]This is a corollary of Steenrod's classification theorem [56].

Taub-NUT configurations are usually considered as the gravitational analog of Dirac's magnetic monopole. Our results show that strictly speaking this is not true. Indeed, Dirac's monopole is treated as a quantum system as a result of the underlying space topology. We have shown that in gravity this role is played by Taub-Bolt configurations, only. We believe that this result might be important for the quantization of gravity by using the path-integral approach. In our opinion, Taub-Bolt instantons should be used to construct the physical vacuum state of gravity. The consequences of this alternative approach are difficult to predict in advance. We expect to handle this problem in a future investigation.

Introducing a cosmological constant to the dyonic instantons does not alter their topology. In fact, our results remain the same if the ALF configurations we have studied are generalized to be asymptotically locally AdS_4. Through the AdS/CFT correspondence[57, 58, 59, 60] dyonic AdS_4 black holes have already been investigated and related to magnetohydrodynamics [61] and Hall conductivity [62]. This points to further exploring the dyonic Taub-NUT/Bolt system.

Acknowledgements

DFA is supported by CONACyT, Grant No. 404449. This work has been supported by the UNAM-DGAPA-PAPIIT, Grant No. IN111617.

References

[1] S.W. Hawking. Gravitational instantons. *Physics Letters A*, 60(2), 1977.

[2] V.A. Belinskii, G.W. Gibbons, D.N. Page, and C.N. Pope. Asymptotically Euclidean Bianchi IX metrics in quantum gravity. *Physics Letters B*, 76(4):433–435, 1978.

[3] G. W. Gibbons and S. W. Hawking. Action integrals and partition functions in quantum gravity. *Physical Review D*, 15(10):2752–2756, 1977.

[4] G. W. Gibbons and S. W. Hawking. Gravitational multi-instantons. *Physics Letters B*, 78(4):430–432, 1978.

[5] Tohru Eguchi and Andrew J. Hanson. Self-dual solutions to Euclidean gravity. *Annals of Physics*, 120:82–106, 1979.

[6] Tohru Eguchi, Peter B. Gilkey, and Andrew J. Hanson. Gravitation, gauge theories and differential geometry. *Physics Reports*, 6(6), 1980.

[7] S. Carlip. Quantum gravity: a progress report. *Reports on Progress in Physics*, 64(8):885–942, 2001.

[8] L. Patiño and H. Quevedo. Topological quantization of gravitational fields. *Journal of Mathematical Physics*, 46(022502), 2005.

[9] F. Nettel and H. Quevedo. Topological spectrum of classical configurations. *AIP Conference Proceedings*, 956(9):9–14, nov 2007.

[10] F. Nettel, H. Quevedo, and M. Rodríguez. Topological spectrum of mechanical systems. *Reports on Mathematical Physics*, 64(3):355–365, dic 2009.

[11] F. Nettel and H. Quevedo. Topological spectrum of the harmonic oscillator. *International Journal of Pure and Applied Mathematics*, 70(2):117–123, 2011.

[12] G. Arciniega, F. Nettel, L. Patiño, and H. Quevedo. Topological quantization of free massive bosonic fields. *Journal of Geometry and Symmetry in Physics*, 26:1–12, 2012.

[13] G. Arciniega, F. Nettel, L. Patiño, and H. Quevedo. Topological discretization of bosonic strings. *International Journal of Pure and Applied Mathematics*, 82(1):65–85, 2013.

[14] P.A.M. Dirac. Quantised singularities in the electromagnetic field. *Proceedings of the Royal Society A*, 133(821):60–72, September 1931.

[15] Gerardus 't Hooft. Magnetic monopoles in unified gauge theories. *Nuclear Physics*, B79:276–284, 1974.

[16] A.M. Polyakov. Particle spectrum in quantum field theory. *Journal of Experimental and Theoretical Physics Letters*, 20(6):194–195, 1974.

[17] A.A. Belavin, A.M. Polyakov, A.S.Schwartz, and Yu.S. Tyupkin. Pseudoparticle solutions of Yang-Mills equations. *Physics Letters*, 59B(1):85–87, 1975.

[18] Tai Tsun Wu and Chen Ning Yang. Concept of nonintegrable phase factors and global formulation of gauge fields. *Physical Review D*, 12(12):3845–3857, 1975.

[19] Ralph M. Kaufmann, Dan Li, and Birgit Wehefritz-Kaufmann. Notes on topological insulators. arxiv: 1501.02874 [math-ph].

[20] C. L. Kane and E. J. Mele. $\mathbb{Z}_2$ topological order and the quantum spin Hall effect. *Physical Review Letters*, 95(146802), 2005.

[21] Liang Fu and C. L. Kane. Time reversal polarization and a $\mathbb{Z}_2$ adiabatic spin pump. *Physical Review B*, 74(195312), 2006.

[22] Liang Fu, C. L. Kane, and E. J. Mele. Topological insulators in three dimensions. *Physical Review Letters*, 98(106803), 2007.

[23] J. E. Moore and L. Balents. Topological invariants of time-reversal-invariant band structures. *Physical Review B*, 75(121306), 2007.

[24] Markus König, Steffen Wiedmann, Christoph Brüne1, Andreas Roth, Hartmut Buhmann, Laurens W. Molenkamp, Xiao-Liang Qi, and Shou-Cheng Zhang. Quantum spin Hall insulator state in HgTe quantum wells. *Science*, 318(5851):766–770, 2007.

[25] M.Z. Hasan and C.L. Kane. Colloquium: Topological insulators. *Review of Modern Physics*, 82:3045–3067, 2010.

[26] Joseph Maciejko, Taylor L. Hughes, , and Shou-Cheng Zhang. The quantum spin Hall effect. *Annual Review of Condensed Matter Physics*, 2:31–53, 2011.

[27] Takahiro Fukui, Ken Shiozaki, Takanori Fujiwara, and Satoshi Fujimoto. Bulk-edge correspondence for Chern topological phases: A viewpoint from a generalized index theorem. *Journal of the Physical Society of Japan*, 81(114602), 2012.

[28] James de Lisle, Suvabrata De, Emilio Alba, Alex Bullivant, Juan J Garcia-Ripoll, Ville Lahtinen, and Jiannis K Pachos. Detection of Chern numbers and entanglement in topological two-species systems through subsystem winding numbers. *New Journal of Physics*, 16(083022), 2014.

[29] D.J. Thouless, M. Kohmoto, M.P. Nightingale, and M. den Nijs. Quantized Hall conductance in a two-dimensional periodic potential. *Physical Review Letters*, 49(6):405–408, 1982.

[30] Barry Simon. Holonomy, the quantum adiabatic theorem, and Berry's phase. *Physical Review Letters*, 51(24):2167–2170, 1983.

[31] M. V. Berry. Quantal phase factors accompanying adiabatic changes. *Proceedings of the Royal Society A*, 392:45–57, 1984.

[32] Joseph E. Avron, Ruedi Seiler, and Barry Simon. The quantum Hall effect and the relative index for projections. *Physical Review Letters*, 65(17):2185–2188, 1990.

[33] J. Bellissard, A. van Elst, and H. Schulz Baldes. The noncommutative geometry of the quantum Hall effect. *Journal of Mathematical Physics*, 35(10):53735451, 1994.

[34] S.W. Hawking and D.N. Page. Thermodynamics of black holes in Anti-de Sitter space. *Communications in Mathematical Physics*, 87:577–588, 1983.

[35] Matthew Headrick and Toby Wiseman. Ricci flow and black holes. *Classical and Quantum Gravity*, 23:6683–6707, 2006.

[36] Clifford V Johnson. The extended thermodynamic phase structure of Taub-NUT and Taub-Bolt. *Class. Quantum Grav.*, 31(22), 2014.

[37] Gustav Holzegel, Thomas Schmelzer, and Claude Warnick. Ricci flows connecting Taub-Bolt and Taub-NUT metrics. *Classical and Quantum Gravity*, 24(24):6201–6217, 2007.

[38] G.W. Gibbons. The Sen conjecture for fundamental monopoles of distinct types. *Physics Letters B*, 382:53–59, 1996.

[39] A. H. Taub. Empty space-times admitting a three parameter group of motions. *Annals of Mathematics*, 53(3):472–490, 1951.

[40] E. Newman, L. Tamburino, and T. Unti. Empty-space generalization of the Schwarzschild metric. *Journal of Mathematical Physics*, 4(7):915–923, 1963.

[41] C. W. Misner. The flatter regions of Newman, Unti and Tamburino's generalized Schwarzschild space. *Journal of Mathematical Physics*, 4(7):924–937, 1963.

[42] H. Stephani, D. Kramer, M. MacCallum, C. Hoenselaers, and E. Herlt. *Exact solutions of Einstein's field equations*. Cambridge University Press, 2009.

[43] Hernando Quevedo. Complex transformations of the curvature tensor. *General Relativity and Gravitation*, 24(7):693–703, 1992.

[44] Dieter R. Brill. Electromagnetic fields in a homogeneous, nonisotropic universe. *Physical Review*, 133(3B):B845–B848, 1964.

[45] G.W. Gibbons and Malcolm J. Perry. New gravitational instantons and their interactions. *Physical Review D*, 22(2):313–321, 1980.

[46] Don N. Page. Taub-NUT instanton with an horizon. *Physics Letters B*, 78(249), 1978.

[47] Andrew Chamblin, Roberto Emparan, Clifford V. Johnson, and Robert C. Myers. Large N phases, gravitational instantons, and the nuts and bolts of AdS holography. *Physical Review D*, 59(064010), 1999.

[48] M. H. Dehghani, S. H. Hendi. Taub-NUT/Bolt Black Holes in Gauss-Bonnet-Maxwell Gravity. *Physical Review D*, 73(084021), 2006.

[49] S. H. Hendi, M. H. Dehghani Taub-NUT Black Holes in Third order Lovelock Gravity *Physics Letters B*, 666:116-120, 2008.

[50] F. Faure and B. Zhilinskii. Topological Chern indices in molecular spectra. *Physical Review Letters*, 85(5):960–963, 2000.

[51] F. Faure and B. Zhilinskii. Topologically coupled energy bands in molecules. *Physics Letters A*, 302:242–252, 2002.

[52] T.H.R. Skyrme. A unified field theory of mesons and baryons. *Nuclear Physics*, 31:556–569, 1962.

[53] C.N. Pope. Axial-vector anomalies and the index theorem in charged Schwarzschild and Taub-NUT space. *Nuclear Physics B*, 141:432–444, 1978.

[54] Gábor Etesi and Tamás Hausel. Geometric interpretation of Schwarzschild instantons. *Journal of Geometry and Physics*, 37:126–136, 2001.

[55] A.N. Aliev and Cihan Saçlıoğu. Self-dual fields harbored by a Kerr-Taub-Bolt instanton. *Physics Letters B*, 632:725–727, 2006.

[56] Norman Steenrod. *The Topology of Fiber Bundles*. Princeton University Press, 1951.

[57] J.M. Maldacena. The large N limit of superconformal field theories and supergravity. *Advances in Theoretical and Mathematical Physics*, 2:231–252, 1998.

[58] S.S. Gubser, I.R. Klebanov, and A.M. Polyakov. Gauge theory correlators from non-critical string theory. *Physics Letters B*, 428:105–114, 1998.

[59] E. Witten. Anti-de Sitter space and holography. *Advances in Theoretical and Mathematical Physics*, 2:253–291, 1998.

[60] E. Witten. Anti-de Sitter space, thermal phase transition, and confinement in gauge theories. *Advances in Theoretical and Mathematical Physics*, 2:505–532, 1998.

[61] Marco M. Caldarelli, Óscar J.C. Dias, and Dietmar Klemm. Dyonic AdS black holes from magnetohydrodynamics. *Journal of High Energy Physics*, 2009(3):025, 2009.

[62] Sean A. Hartnoll and Pavel K. Kovtun. Hall conductivity from dyonic black holes. *Physical Review D*, 76(6), 2007.